

A uniqueness theorem for meromorphic functions ignoring multiplicity

Nguyen Duy Phuong (Thai Nguyen, Vietnam)

(Received Mar. 13, 2023)

Abstract. In this paper, we give a uniqueness theorem for meromorphic functions ignoring multiplicity, which generalizes a An's theorem in [1].

1. Introduction. Main results

In this paper, by a meromorphic function we mean a meromorphic function on the complex plane \mathbb{C} .

Let f be a non-constant meromorphic function on \mathbb{C} . For every $a \in \mathbb{C}$, we define the function $\nu_f^a : \mathbb{C} \rightarrow \mathbb{N}$ by

$$\nu_f^a(z) = \begin{cases} 0 & \text{if } f(z) \neq a \\ d & \text{if } f(z) = a \text{ with multiplicity } d, \end{cases}$$

and set $\nu_f^\infty = \nu_{\frac{1}{f}}^0$, and define the function $\bar{\nu}_f^a : \mathbb{C} \rightarrow \mathbb{N}$ by $\bar{\nu}_f^a(z) = \min \{ \nu_f^a(z), 1 \}$, and set $\bar{\nu}_f^\infty = \bar{\nu}_{\frac{1}{f}}^0$. For $f \in \mathcal{M}(\mathbb{C})$ and a non-empty set $S \subset \mathbb{C} \cup \{\infty\}$, we define

$$E_f(S) = \bigcup_{a \in S} \{ (z, \nu_f^a(z)) : z \in \mathbb{C} \}, \quad \bar{E}_f(S) = \bigcup_{a \in S} \{ (z, \bar{\nu}_f^a(z)) : z \in \mathbb{C} \}.$$

Let \mathcal{F} be a nonempty subset of $\mathcal{M}(\mathbb{C})$. Two functions f, g of \mathcal{F} are said to *share S , counting multiplicity* (share S CM) if $E_f(S) = E_g(S)$, and to *share S , ignoring multiplicity* (share S IM) if $\bar{E}_f(S) = \bar{E}_g(S)$.

Key words and phrases: Meromorphic Function, uniqueness, ignoring multiplicity.
2020 Mathematics Subject Classification: 30D05

If the condition $E_f(S) = E_g(S)$ implies $f = g$ for any two non-constant meromorphic (entire) functions f, g , then S is called a unique range set for meromorphic (entire) functions counting multiplicity, or in brief, URSM (URSE). A set $S \subset \mathbb{C} \cup \{\infty\}$ is called a unique range set for meromorphic (entire) functions ignoring multiplicity, or in brief, URSM-IM (URSE-IM), if the condition $\overline{E}_f(S) = \overline{E}_g(S)$ implies $f = g$ for any pair of non-constant meromorphic (entire) functions.

In 1976 Gross ([10]) proved that there exist three finite sets S_j ($j = 1, 2, 3$) such that any two entire functions f and g satisfying $E_f(S_j) = E_g(S_j)$, $j = 1, 2, 3$ must be identical. In the same paper Gross([10]) posed the following question:

Question A. *Can one find two (or possible even one) finite set S_j ($j = 1, 2$) such that any two entire functions f and g satisfying $E_f(S_j) = E_g(S_j)$ ($j = 1, 2$) must be identical?*

Yi ([18]-[20],[22]) first gave an affirmative answer to Question A. Since then, many results have been obtained for this and related topics (see ([1]-[15]), ([17]-[23])).

Concerning to Question A, a natural question is the following.

Question B. *What is the smallest cardinality for such a finite set S such that any two meromorphic functions f and g satisfying either $E_f(S) = E_g(S)$ or $\overline{E}_f(S) = \overline{E}_g(S)$ must be identical?*

So far, the best answer to Question B for the case of URSM was obtained by Frank and Reinders ([7]). They proved the following result.

Theorem C. *The set $\{z \in \mathbb{C} \mid P_{FR}(z) = \frac{(n-1)(n-2)}{2}z^n + n(n-2)z^{n-1} + \frac{(n-1)n}{2}z^{n-2} - c = 0\}$, where $n \geq 11$ and $c \neq 0, 1$, is a unique range set for meromorphic functions counting multiplicity.*

In 1997, H. X. Yi ([21]) first gave an answer to question B for the case of URSM-IM with 19 elements. Since then, many results have been obtained for this topic (see ([1]- [5])). So far, the best answer to Question B for the case of URSM-IM was obtained by Chakraborty([5]). He proved the following result.

Theorem D. *Let $S_{FR} = \{z \in \mathbb{C} \mid P_{FR}(z) = 0\}$. If $n \geq 15$, then S_{FR} is a URSM-IM.*

In 2022, An([1]) given a class of unique range sets for meromorphic functions ignoring multiplicity with 15 elements. He proved the following result.

Let $n \in \mathbb{N}^*$, $n \geq 3$. Consider polynomial $P(z)$ as follows:

$$P_A(z) = z^n - \frac{2na}{n-1}z^{n-1} + \frac{na^2}{n-2}z^{n-2} + 1 = Q_A(z) + 1, \quad (1.1)$$

where $a \in \mathbb{C}$, $a \neq 0$. Suppose that

$$Q_A(a) \neq -1, \quad (1.2)$$

$$Q_A(a) \neq -2. \quad (1.3)$$

Theorem E. *Let $P_A(z)$ be defined by (1.1) with conditions (1.2) and (1.3), and let $S_A = \{z \in \mathbb{C} \mid P_A(z) = 0\}$. If $n \geq 15$, then S_A is a URSM-IM.*

Clearly, $P'_A(z) = nz^{n-3}(z-a)^2$, and $P'_{FR}(z) = \frac{n(n-1)(n-2)}{2}z^{n-3}(z-1)^2$. Therefore, this class is different from Chakraborty's Theorem D in ([5]).

In this paper, we give a uniqueness theorem for meromorphic functions ignoring multiplicity, which generalizes Theorem E.

Now let us describe main results of the paper.

Let $q, k, m_1, m_2 \in \mathbb{N}^*$.

We will let $P(z)$ be polynomial having no multiple zeros of degree q in $\mathbb{C}[z]$:

$$P(z) = (m_1 + m_2 + 1) \left(\sum_{i=0}^{m_2} \binom{m_2}{i} \frac{(-1)^i}{m_1 + m_2 + 1 - i} z^{m_1 + m_2 + 1 - i} a^i \right) + 1 = Q(z) + 1,$$

where

$$Q(z) = (m_1 + m_2 + 1) \left(\sum_{i=0}^{m_2} \binom{m_2}{i} \frac{(-1)^i}{m_1 + m_2 + 1 - i} z^{m_1 + m_2 + 1 - i} a^i \right). \quad (1.4)$$

Suppose that

$$a \neq 0, Q(a) \neq -1, Q(a) \neq -2. \quad (1.5)$$

Clearly, $P'(z) = (m_1 + m_2 + 1)z^{m_1}(z-a)^{m_2}$, and has a zero at 0 of order m_1 , and a zero at a of order m_2 . Note that $q = m_1 + m_2 + 1$.

We shall prove the following theorem.

Theorem 1. *Let $P(z)$ be defined in (1.4) with conditions (1.5), and let $S = \{z \in \mathbb{C} \mid P(z) = 0\}$. If $q \geq 15$, then S is a URSM-IM.*

Remark 2. *From proof of Theorem 1 we give a proof of Theorem D, which is different from Chakraborty's proof in ([5]) (see section 3.).*

Remark 3. *In Theorem 1, take $m_1 = n - 3$ and $m_2 = 2$ we obtain Theorem E.*

Indeed, by $P'_A(z) = nz^{n-3}(z-a)^2$ and $P'(z) = (m_1 + m_2 + 1)z^{m_1}(z-a)^{m_2}$, we obtain $P(z) = P_A(z)$ when $m_1 = n - 3$, $m_2 = 2$.

2. Lemmas, Definitions

We assume that the reader is familiar with the notations of Nevanlinna theory (see, for example, ([6]), ([16])). We need some lemmas.

Lemma 2.1. ([6], paper 98; [16], paper 43) *Let f be a non-constant meromorphic function on \mathbb{C} and let a_1, a_2, \dots, a_q be distinct points of $\mathbb{C} \cup \{\infty\}$. Then*

$$(q-2)T(r, f) \leq \sum_{i=1}^q \overline{N}\left(r, \frac{1}{f-a_i}\right) - N_0\left(r, \frac{1}{f'}\right) + S(r, f),$$

where $N_0(r, \frac{1}{f'})$ is the counting function of those zeros of f' , which are not zeros of function $(f-a_1)\dots(f-a_q)$, and $S(r, f) = o(T(r, f))$ for all r , except for a set of finite Lebesgue measure.

Lemma 2.2. ([6, paper 99]) *For any non-constant meromorphic function f ,*

$$T\left(r, \frac{1}{f'}\right) \leq 2T(r, f) + S(r, f).$$

Definition. Let f be a non-constant meromorphic function, and k be a positive integer. We denote by $\overline{N}_{(k)}(r, f)$ the counting function of the poles of order $\geq k$ of f , where each pole is counted only once. If z is a zero of f , denote by $\nu_f(z)$ its multiplicity. We denote by $\overline{N}(r, \frac{1}{f'}; f \neq 0)$ the counting function of the zeros z of f' satisfying $f(z) \neq 0$, where each zero is counted only once.

Let be given two non-constant meromorphic functions f and g . For simplicity, denote by $\nu_1(z) = \nu_f(z)$ (resp., $\nu_2(z) = \nu_g(z)$), if z is a zero of f (resp., g). Let $f^{-1}(0) = g^{-1}(0)$. We denote by $N(r, \frac{1}{f}; \nu_1 = \nu_2 = 1)$ (resp., $\overline{N}(r, \frac{1}{f}; \nu_1 > \nu_2 \geq 1)$) the counting function of the common zeros z , satisfying $\nu_1(z) = \nu_2(z) = 1$ (resp., $\nu_1(z) > \nu_2(z) \geq 1$, where each zero is counted only once), and by $N(r, \frac{1}{f}; \nu_1 \geq 2)$ the counting function of the zeros z of f , satisfying $\nu_1(z) \geq 2$. Similarly, we define the counting functions $\overline{N}(r, \frac{1}{g}; \nu_2 > \nu_1 \geq 1)$, $N(r, \frac{1}{g}; \nu_2 \geq 2)$.

Lemma 2.3. ([1, Lemma 2.3])

Let f, g be two non-constant meromorphic functions and let $f^{-1}(0) = g^{-1}(0)$. Set

$$F = \frac{1}{f}, G = \frac{1}{g}, L = \frac{F''}{F'} - \frac{G''}{G'}.$$

Suppose that $L \neq 0$. Then

$$\begin{aligned} 1) \quad N(r, L) &\leq \overline{N}_{(2)}(r, f) + \overline{N}_{(2)}(r, g) + \\ &\overline{N}(r, \frac{1}{f}; \nu_1 > \nu_2 \geq 1) + \overline{N}(r, \frac{1}{g}; \nu_2 > \nu_1 \geq 1) + \overline{N}(r, \frac{1}{f}; f \neq 0) + \\ &\overline{N}(r, \frac{1}{g}; g \neq 0). \end{aligned}$$

Moreover, if a is a common simple zero of f and g , then $L(a) = 0$.

$$\begin{aligned} 2) \quad \overline{N}(r, \frac{1}{f}) + \overline{N}(r, \frac{1}{g}) + \overline{N}(r, \frac{1}{f}; \nu_1 > \nu_2 \geq 1) + \overline{N}(r, \frac{1}{g}; \nu_2 > \nu_1 \geq 1) \\ \leq N(r, L) + \frac{1}{2}(N(r, \frac{1}{f}) + N(r, \frac{1}{g})) + N(r, \frac{1}{f}; \nu_1 \geq 2) + N(r, \frac{1}{g}; \nu_2 \geq 2) \\ + S(r, f) + S(r, g). \end{aligned}$$

A polynomial $R(z)$ is called a *strong uniqueness polynomial for meromorphic (entire) functions* if for arbitrary two non-constant meromorphic (entire) functions f and g , and a nonzero constant c , the condition $R(f) = cR(g)$ implies $f = g$ (see ([2]), ([9]), ([13])). In this case we say $R(z)$ is a SUPM (SUPE). A polynomial $R(z)$ is called a *uniqueness polynomial for meromorphic (entire) functions* if for arbitrary two non-constant meromorphic (entire) functions f and g , the condition $R(f) = R(g)$ implies $f = g$ (see ([2]), ([9]), ([13])). In this case we say $R(z)$ is a UPM (UPE). Let $R(z)$ be a polynomial of the degree q . Assume that the derivative of $R(z)$ has mutually distinct k zeros d_1, d_2, \dots, d_k with multiplicities q_1, q_2, \dots, q_k , respectively. We often consider polynomials satisfying the following condition introduced by Fujimoto ([8]):

$$R(d_i) \neq R(d_j), 1 \leq i < j \leq k. \quad (2.1)$$

The number k is called the *derivative index* of R .

H. Fujimoto ([8], Proposition 7.1)) proved the following:

Lemma 2.4. *Let $R(z)$ be a polynomial of degree q satisfying the condition (2.1), we assume furthermore that $q \geq 5$ and there are two non-constant meromorphic function f and g such that*

$$\frac{1}{R(f)} = \frac{c_0}{R(g)} + c_1$$

for two constants $c_0 \neq 0$ and c_1 . If $k \geq 3$ or if $k = 2, \min\{q_1, q_2\} \geq 2$, then $c_1 = 0$.

Lemma 2.5. ([13], Theorem 1.1)

Let $P(z)$ be defined by (1.4) with conditions (1.5), and let $n \geq 6$. Then $P(z)$ is a strong uniqueness polynomial for meromorphic functions.

Lemma 2.6. ([3], Theorem 1.1)

Let $P_{FR_1}(z) = \frac{(n-1)(n-2)}{2}z^n + n(n-2)z^{n-1} + \frac{(n-1)n}{2}z^{n-2} - c = 0$, where $n \geq 8$ and $c \in \mathbb{C}$. Then $P_{FR_1}(z)$ is a strong uniqueness polynomial for meromorphic functions.

3. Proof of Theorems**Proof of Theorem 1**

Recall that $P(z) = (z - a_1) \dots (z - a_q)$, $P'(z) = qz^{m_1}(z - a)^{m_2}$, $q = m_1 + m_2 + 1$.

Suppose $q \geq 15$ and $\overline{E}_f(S) = \overline{E}_g(S)$, where $S = \{z \in \mathbb{C} \mid P(z) = 0\}$. Set

$$F = \frac{1}{P(f)}, G = \frac{1}{P(g)}, L = \frac{F''}{F'} - \frac{G''}{G'},$$

$$T(r) = T(r, f) + T(r, g), S(r) = S(r, f) + S(r, g).$$

Then $T(r, P(f)) = qT(r, f) + S(r, f)$ and $T(r, P(g)) = qT(r, g) + S(r, g)$, and hence $S(r, P(f)) = S(r, f)$ and $S(r, P(g)) = S(r, g)$.

We consider two following cases:

Case 1. $L \equiv 0$. Then, we have $\frac{1}{P(f)} = \frac{c}{P(g)} + c_1$ for some constants $c \neq 0$ and c_1 . By Lemma 2.4 we obtain $c_1 = 0$.

Therefore, there is a constant $C \neq 0$ such that $P(f) = CP(g)$. Then, applying Lemma 2.5 we obtain $f = g$.

Case 2. $L \not\equiv 0$.

Claim 1. We have

$$(q-2)T(r) \leq \overline{N}(r, \frac{1}{P(f)}) + \overline{N}(r, \frac{1}{P(g)}) - N_0(r, \frac{1}{f'}) - N_0(r, \frac{1}{g'}) + S(r), \quad (3.1)$$

where $N_0(r, \frac{1}{f'})$ ($N_0(r, \frac{1}{g'})$) is the counting function of those zeros of f' , which are not zeros of function $(f - a_1) \dots (f - a_q)f(f - a)((g - a_1) \dots (g - a_q)g(g - a))$.

Indeed, applying the Lemma 2.1 to the functions f, g and the values $a_1, a_2, \dots, a_q, 0, a, \infty$, and noting that

$$\sum_{i=1}^q \overline{N}\left(r, \frac{1}{f - a_i}\right) = \overline{N}\left(r, \frac{1}{P(f)}\right), \quad \sum_{i=1}^q \overline{N}\left(r, \frac{1}{g - a_i}\right) = \overline{N}\left(r, \frac{1}{P(g)}\right),$$

we obtain

$$(q+1)T(r) \leq \overline{N}(r, f) + \overline{N}(r, g) + \overline{N}\left(r, \frac{1}{P(f)}\right) + \overline{N}\left(r, \frac{1}{P(g)}\right) + \overline{N}\left(r, \frac{1}{f}\right) + \overline{N}\left(r, \frac{1}{g}\right) + \overline{N}\left(r, \frac{1}{f-a}\right) + \overline{N}\left(r, \frac{1}{g-a}\right) - N_0\left(r, \frac{1}{f'}\right) - N_0\left(r, \frac{1}{g'}\right) + S(r). \quad (3.2)$$

On the other hand,

$$\begin{aligned} \overline{N}(r, f) + \overline{N}(r, g) &\leq (T(r, f) + T(r, g)) + S(r) = T(r) + S(r), \\ \overline{N}\left(r, \frac{1}{f}\right) + \overline{N}\left(r, \frac{1}{g}\right) &\leq (T(r, f) + T(r, g)) + S(r) = T(r) + S(r), \\ \overline{N}\left(r, \frac{1}{f-a}\right) + \overline{N}\left(r, \frac{1}{g-a}\right) &\leq (T(r, f) + T(r, g)) + S(r) = T(r) + S(r). \end{aligned}$$

From this and (3.2) we obtain (3.1).

Claim 2. We have

$$\begin{aligned} \overline{N}\left(r, \frac{1}{P(f)}\right) + \overline{N}\left(r, \frac{1}{P(g)}\right) &\leq \\ \left(\frac{q}{2} + 3\right)T(r) + \overline{N}\left(r, \frac{1}{[P(f)]'}; P(f) \neq 0\right) + \overline{N}\left(r, \frac{1}{[P(g)]'}; P(g) \neq 0\right) + S(r). \end{aligned}$$

Indeed, by $\overline{E}_f(S) = \overline{E}_g(S)$ we get $(P(f))^{-1}(0) = (P(g))^{-1}(0)$. For simplicity, we set $\nu_1 = \nu_1(z)$, $\nu_2 = \nu_2(z)$, where $\nu_1(z) = \nu_{P(f)}(z)$, $\nu_2(z) = \nu_{P(g)}(z)$. Note that

$$\begin{aligned} \overline{N}_{(2)}(r, P(f)) &= \overline{N}(r, f), \quad \overline{N}_{(2)}(r, P(g)) = \overline{N}(r, g), \\ S(r, P(f)) &= S(r, f), \quad S(r, P(g)) = S(r, g), \quad S(r) = S(r, f) + S(r, g). \end{aligned}$$

Applying the Lemma 2.3 to the functions $P(f), P(g)$. Then we obtain

$$\begin{aligned} N(r, L) &\leq \overline{N}(r, f) + \overline{N}(r, g) + \overline{N}\left(r, \frac{1}{P(f)}; \nu_1 > \nu_2 \geq 1\right) + \overline{N}\left(r, \frac{1}{P(g)}; \nu_2 > \nu_1 \geq 1\right) \\ &\quad + \overline{N}\left(r, \frac{1}{[P(f)]'}; P(f) \neq 0\right) + \overline{N}\left(r, \frac{1}{[P(g)]'}; P(g) \neq 0\right), \end{aligned} \quad (3.3)$$

and

$$\begin{aligned} & \overline{N}(r, \frac{1}{P(f)}) + \overline{N}(r, \frac{1}{P(g)}) + \overline{N}(r, \frac{1}{P(f)}; \nu_1 > \nu_2 \geq 1) + \\ & \overline{N}(r, \frac{1}{P(g)}; \nu_2 > \nu_1 \geq 1) \leq N(r, L) + \frac{1}{2}(N(r, \frac{1}{P(f)}) + N(r, \frac{1}{P(g)})) + \\ & N(r, \frac{1}{P(f)}; \nu_1 \geq 2) + N(r, \frac{1}{P(g)}; \nu_2 \geq 2) + S(r). \end{aligned} \quad (3.4)$$

Moreover,

$$\overline{N}(r, f) + \overline{N}(r, g) \leq T(r) + S(r). \quad (3.5)$$

Obviously,

$$\begin{aligned} N(r, \frac{1}{P(f)}) & \leq qT(r, f) + S(r, f); N(r, \frac{1}{P(g)}) \leq qT(r, g) + S(r, g), \\ N(r, \frac{1}{P(f)}) + N(r, \frac{1}{P(g)}) & \leq qT(r) + S(r). \end{aligned} \quad (3.6)$$

On the other hand, from $P(f) = (f - a_1) \dots (f - a_q)$ it follows that if z_0 zero is a zero of $P(f)$ with multiplicity ≥ 2 , then z_0 is a zero of $f - a_i$ with multiplicity ≥ 2 for some $i \in \{1, 2, \dots, q\}$, and therefore, it is a zero of f' , so we have

$$N(r, \frac{1}{P(f)}; \nu_1 \geq 2) \leq N(r, \frac{1}{f'}).$$

From this and Lemma 2.2 we obtain

$$N(r, \frac{1}{P(f)}; \nu_1 \geq 2) \leq N(r, \frac{1}{f'}) \leq T(r, f') + S(r, f) \leq 2T(r, f) + S(r, f).$$

Similarly, we have

$$N(r, \frac{1}{P(g)}; \nu_2 \geq 2) \leq N(r, \frac{1}{g'}) \leq T(r, g') + S(r, g) \leq 2T(r, g) + S(r, g).$$

Therefore,

$$N(r, \frac{1}{P(f)}; \nu_1 \geq 2) + N(r, \frac{1}{P(g)}; \nu_2 \geq 2) \leq 2T(r) + S(r). \quad (3.7)$$

Combining (3.1)-(3.7) we get

$$\begin{aligned} & \overline{N}(r, \frac{1}{P(f)}) + \overline{N}(r, \frac{1}{P(g)}) \leq \\ & (\frac{q}{2} + 3)T(r) + \overline{N}(r, \frac{1}{[P(f)]'}; P(f) \neq 0) + \overline{N}(r, \frac{1}{[P(g)]'}; P(g) \neq 0) + S(r). \end{aligned}$$

Claim 2 is proved.

Claim 3. We have

$$\begin{aligned} \overline{N}(r, \frac{1}{[P(f)]'}; P(f) \neq 0) + \overline{N}(r, \frac{1}{[P(g)]'}; P(g) \neq 0) &\leq 2T(r) + N_0(r, \frac{1}{f'}) + \\ &N_0(r, \frac{1}{g'}) + S(r). \end{aligned}$$

We have

$$\begin{aligned} \overline{N}(r, \frac{1}{[P(f)]'}; P(f) \neq 0) &= \overline{N}(r, \frac{1}{f^{m_1}(f-a)^{m_2}f'}; P(f) \neq 0) \leq \overline{N}(r, \frac{1}{f}) + \\ \overline{N}(r, \frac{1}{f-a}) + \overline{N}_0(r, \frac{1}{f'}) &\leq 2T(r, f) + \overline{N}_0(r, \frac{1}{f'}) + S(r, f). \end{aligned} \quad (3.8)$$

Similarly,

$$\overline{N}(r, \frac{1}{[P(g)]'}; P(g) \neq 0) \leq 2T(r, g) + \overline{N}_0(r, \frac{1}{g'}) + S(r, g). \quad (3.9)$$

Inequalities (3.8) and (3.9) give us

$$\begin{aligned} \overline{N}(r, \frac{1}{[P(f)]'}; P(f) \neq 0) + \overline{N}(r, \frac{1}{[P(g)]'}; P(g) \neq 0) &\leq \\ &\leq 2T(r) + \overline{N}_0(r, \frac{1}{f'}) + \overline{N}_0(r, \frac{1}{g'}) + S(r). \end{aligned}$$

Claim 3 is proved.

Claim 1, 2, 3 give us:

$$(q-2)T(r) \leq (\frac{q}{2} + 5)T(r) + S(r). \text{ So } (q-14)T(r) \leq S(r).$$

This is a contradiction to the assumption that $q \geq 15$. So $L \equiv 0$. Therefore $f = g$. Theorem 1 is proved.

A proof of Theorem D

By using the arguments similar in proof of Theorem 1 and Lemma 2.6 we give a proof of Theorem D, which is different from Chakraborty's proof of Theorem D in ([5]).

Recall that $P_{FR}(z) = (z-a_1)\dots(z-a_n)$, $P'_{FR}(z) = \frac{n(n-1)(n-2)}{2} z^{n-3}(z-1)^2$.

Suppose $n \geq 15$ and $\overline{E}_f(S_{FR}) = \overline{E}_g(S_{FR})$, where $S_{FR} = \{z \in \mathbb{C} \mid P_{FR}(z) = 0\}$. Set

$$F = \frac{1}{P_{FR}(f)}, G = \frac{1}{P_{FR}(g)}, L = \frac{F''}{F'} - \frac{G''}{G'},$$

$$T(r) = T(r, f) + T(r, g), S(r) = S(r, f) + S(r, g).$$

Then $T(r, P_{FR}(f)) = nT(r, f) + S(r, f)$ and $T(r, P_{FR}(g)) = nT(r, g) + S(r, g)$, and hence $S(r, P_{FR}(f)) = S(r, f)$ and $S(r, P_{FR}(g)) = S(r, g)$.

We consider two following cases:

Case 1. $L \equiv 0$. Then, we have $\frac{1}{P_{FR}(f)} = \frac{c}{P_{FR}(g)} + c_1$ for some constants $c \neq 0$ and c_1 . By Lemma 2.4 we obtain $c_1 = 0$.

Therefore, there is a constant $C \neq 0$ such that $P_{FR}(f) = CP_{FR}(g)$. Then, applying Lemma 2.6 we obtain $f = g$.

Case 2. $L \not\equiv 0$. By using the arguments similar in proof of Theorem 1 we obtain

Claim 1. We have

$$(n-2)T(r) \leq \overline{N}(r, \frac{1}{P_{FR}(f)}) + \overline{N}(r, \frac{1}{P_{FR}(g)}) - N_0(r, \frac{1}{f'}) - N_0(r, \frac{1}{g'}) + S(r), \quad (3.10)$$

where $N_0(r, \frac{1}{f'})$ ($N_0(r, \frac{1}{g'})$) is the counting function of those zeros of f' , which are not zeros of function $(f - a_1) \dots (f - a_n)f(f-1)((g - a_1) \dots (g - a_n)g(g-1))$.

Claim 2. We have

$$\overline{N}(r, \frac{1}{P_{FR}(f)}) + \overline{N}(r, \frac{1}{P_{FR}(g)}) \leq$$

$$(\frac{n}{2} + 3)T(r) + \overline{N}(r, \frac{1}{[P_{FR}(f)]'}; P_{FR}(f) \neq 0) + \overline{N}(r, \frac{1}{[P_{FR}(g)]'}; P_{FR}(g) \neq 0) + S(r).$$

Claim 3. We have

$$\overline{N}(r, \frac{1}{[P_{FR}(f)]'}; P_{FR}(f) \neq 0) + \overline{N}(r, \frac{1}{[P_{FR}(g)]'}; P_{FR}(g) \neq 0) \leq 2T(r) + N_0(r, \frac{1}{f'}) +$$

$$N_0(r, \frac{1}{g'}) + S(r).$$

Claim 1, 2, 3 give us:

$$(n-2)T(r) \leq (\frac{n}{2} + 5)T(r) + S(r). \text{ So } (n-14)T(r) \leq S(r).$$

This is a contradiction to the assumption that $n \geq 15$. So $L \equiv 0$. Therefore $f = g$.

Theorem D is proved.

References

- [1] Vu Hoai An, *A new class of unique range sets for meromorphic functions ignoring multiplicity with 15 elements*, Journal of Mathematics and Mathematical Sciences(2022), Vol. 1, 1-14.
- [2] A. Banerjee, *A new class of strong uniqueness polynomial satisfying Fujimoto's conditions*, Ann. Acad. Sci. Fenn. Math. Vol. 40, 2015, 465-474.
- [3] A. Banerjee, B. Chakraborty, S. Mallick, *Further Investigations on Fujimoto Type Strong Uniqueness Polynomials*, Filomat 31:16 (2017), 5203-5216.
- [4] S. Bartels, *Meromorphic functions sharing a set with 17 elements ignoring multiplicities*, Compl. Var. Theory Appl., 39, 85-92 (1999).
- [5] B. Chakraborty, *On the Cardinality of a Reduced Unique-Range Set*, Ukr. Math. J., Vol. 72, No. 11, April, 2021, DOI 10.1007/s11253-021-01889-z.
- [6] A. A. Goldberg and I. V. Ostrovskii, *Value Distribution of Meromorphic Functions*, Translations of Mathematical Monographs (2008), V.236.
- [7] G. Frank and M. Reinders, *A unique range set for meromorphic functions with 11 elements*, Compl. Var. Theory Appl. 37:1, 1998, 185-193.
- [8] H. Fujimoto, *On uniqueness of meromorphic functions sharing finite sets*, Amer. J. Math. 122, 2000, 1175-1203.
- [9] H. Fujimoto, *On uniqueness polynomials for meromorphic functions*, Nagoya Math. J., 170, 33-46 (2003).
- [10] F. Gross, *Factorization of meromorphic functions and some open problems*, Complex Analysis(Proc. Conf. Univ. Kentucky, Lexington, Ky. 1976), pp. 51-69, Lecture Notes in Math. Vol. 599, Springer, Berlin, 1977.
- [11] Ha Huy Khoai, *Some remarks on the genericity of unique range sets for meromorphic functions*, Sci. China Ser. A Mathematics, Vol. 48, 2005, 262-267.
- [12] Ha Huy Khoai, Vu Hoai An, and Pham Ngoc Hoa, *On functional equations for meromorphic functions and applications*, Arch. Math, DOI 10.1007/s00013-017-1093-5, 2017.

- [13] Ha Huy Khoai, Vu Hoai An and Nguyen Xuan Lai, *Strong uniqueness polynomials of degree 6 and unique range sets for powers of meromorphic functions*, Intern. J. Math., 2018, DOI:10.1142/S0129167X18500374.
- [14] Ha Huy Khoai, Vu Hoai An and Le Quang Ninh, *Value-sharing and uniqueness for L -functions*, Ann. Polonici Math., 2021, 265-278.
- [15] Ha Huy Khoai and Vu Hoai An, *Determining an L -function in the extended Selberg class by its preimages of subsets*, Ramanujan Journal, 58, 253-267 (2022).
- [16] W.K.Hayman, *Meromorphic Functions*, Clarendon, Oxford(1964).
- [17] P.Li and C.C. Yang, *Some further results on the unique range sets of meromorphic functions*, Kodai Math. J. 18, 1995, 437-450.
- [18] H.X.Yi , *Uniqueness of Meromorphic Functions and question of Gross*, Sci. China (Ser. A), Vol.37 N0.7, July 1994, 802-813.
- [19] H. X. Yi, *A question of Gross and the uniqueness of entire functions*, Nagoya Math. J. Vol. 138 (1995), 169-177.
- [20] H. X. Yi, *Unicity theorems for meromorphic and entire functions III*, Bull. Austral. Math. Soc., 53, 71-82 (1996).
- [21] H. X. Yi, *The reduced unique range sets for entire or meromorphic functions*, Compl. Var. Theory Appl., 32, 191-198 (1997).
- [22] H. X. Yi, *On a question of Gross concerning uniqueness of entire functions*, Bull. Austral. Math. Soc. Vol. 57(1998), 343-349.
- [23] H. X. Yi and W.C.Lin, *Uniqueness theorems concerning a question of Gross*, Proc. Japan Acad., Ser. A, 80, 2004, 136-140.

Nguyen Duy Phuong

Thai Nguyen University of Education

Thai Nguyen

Vietnam

phuongnd@tnu.edu.vn